arXiv:1206.1923v1 [hep-ph] 9 Jun 2012

A proton-pentaquark mixing and the intrinsic charm model

Mikhail Mikhasenko
Iistitute for High Energy Physics, Protvito, Russia

A new interpretation of intrinsic charm phenomenon based on the assumption of pentaquark $|uudc\bar{c}\rangle$ mixing with a proton is offered. The structure function of the c-quark in the pentaquark is constructed. Mixing evaluation is given according to theoretical point of view, experiment data on D-meson production and inclusive production of the hidden charm particles.

Today there are many articles related to intrinsic charm concept, which was introduced by Brodsky et al. [1]. At the current article we want to discuss a new exited interpretation of the intrinsic charm phenomenon. As will be shown below, the intrinsic charm problem is closely related to the pentaquark $|uudc\bar{c}\rangle$ existance and mixing of that state with the proton.

Authors of ref. [1] made the assumption that Fock state decomposition of the proton wave-function contains a non-negligible $|uudc\bar{c}\rangle$ component, which results in the specific distribution of c-quark in proton. According to [1], the probability to find $|uudc\bar{c}\rangle$ configuration, in classical pertrubation theory, is get by expression:

$$W(A \to q_1 \dots q_5) \sim \left| \frac{\langle q_1 \dots q_5 | M | A \rangle}{E_A - E_{q_1} - \dots - E_{q_5}} \right|^2 \tag{1}$$

Here and in the following we neglict any momentum dependence of the numenator $\langle q_1 \dots q_5 | M | A \rangle^2$. This matrix element determs absolute normalisation, which will be discussed below. In the infinite-momentum frame, the energy denominator can be simply experssed in terms of the masses and momuntum fraction x_i of the q_i. Thus for that case $p \to |uudc\bar{c}\rangle$ we have

$$W(p \to |uudc\bar{c}\rangle) \sim \left[m_p^2 - \sum_{i=1}^{5} \frac{m_\perp^2}{x_i} \right]^{-2} \tag{2}$$

Here $m_{i\perp} = \sqrt{m_i^2 + p_\perp^2}$ is a transverse quark mass, m_p is a mass of the proton. It is sufficient to evaluate the distribution in the limit of very heavy quarks: $m_c^2 \gg m_i^2,\ m_p^2$

$$W(p \to |uudc\bar{c}\rangle) \sim \frac{x_4^2 x_5^2}{(x_4 + x_5)^2}, \tag{3}$$

Then authors of [1] get structure function of quarks:

$$dP \sim \delta(1 - \sum_{1}^{5} x_i) \frac{x_4^2 x_5^2}{(x_4 + x_5)^2} \prod_{i=1}^{5} dx_i \tag{4}$$

Repeating the integration over $dx_1 \dots dx_4$ as ref. [1], one can easily obtain the destribution of c-quark:

$$P(x) = N x^2 \left[(1-x)(1 + 10x + x^2) + 6x(1+x) \ln \frac{1}{x} \right], \tag{5}$$

where $N = 600$ is determined by normalization condition for one c-quark $\int P(x)dx = 1$.

The distribution $P(x)$ is shown in fig. 1 by solid line. Under (5) we have the average of one charm quark momentum to be $\frac{2}{7}$, so the intrinsic charm contribution should stand out in D-meson spectrum for $x = \frac{2}{7}$, as well as for an inclusive hidden charm particle production $(J/\Psi, \chi_c)$. It was discussed in ref. [2, 3].

Let us consider alternative model [4]. We construct stucture function for c-quark in the pentaquark $|uudc\bar{c}\rangle$, which has quantum numbers like a proton. The result is obtained with assumption about the Regge asymptotics at $x \to 0$ and the limitation of phase space with the transterse momentum:

$$dP_v(x) \sim \frac{\Psi(x)dx}{\sqrt{x^2 + \frac{\mu^2}{P^2}}}, \tag{6}$$

$$\text{where } \Psi(x) \sim \begin{cases} x^{1-\alpha_i} & \text{for valence quarks with fravor } i, \\ 1 & \text{for sea quarks,} \end{cases}$$

and α_i - the intercept for the leading Regge trajectory related to quark with flavor i.

The distribution function for the n-quarks state with m valence quarks can be drawn, according to [4]:

$$dG(x_1, \dots, x_n) \sim \delta(1 - \sum_{1}^{5} x_i) \prod_{i=1}^{m} x_i^{1-\alpha_i} \prod_{j=1}^{n} \frac{dx_j}{\sqrt{x_j^2 + \frac{\mu^2}{P^2}}} \tag{7}$$

Repeating the calculations of ref. [4] for our case, with diffrent α_i, we get structure function for c-quark in the pentaquark, which consists of $uudc\bar{c}$ quarks:

$$G(x) = M x^{-\alpha_5} (1-x)^{-1+\gamma_A + \sum_1^4 (1-\alpha_i)}, \tag{8}$$

where M is a parameter, which stands unitary normalisation: $\int G(x)dx = 1$.

Values α_i were determed as following: $\alpha_{1,2,3} = \alpha_{u,d}(0) \approx \frac{1}{2}$ are the intersepts for leading Regge f, A_2 trajectories, $\alpha_{4,5} = \alpha_c(0) \approx -2.2$ is the intersept for J/Ψ trajectory, getting at the alticle [5].

In original version Kuti-Weisskopf model [4] the unknown parameter $\gamma_A = 3$ determining the sea normalisation can be obtain using Drell-Yan-West relation [6]. We determ $\gamma_A = 3$ according to [4]:

$$G(x) = Mx^{2.2}(1-x)^{6.7} \tag{9}$$

The structure function of the c-quark for the pentaquark $|uudc\bar{c}\rangle$ (9) is shown in Fig. 1 by dotted line in comparison with the formula (5).



Figure 1: Momentum distrubution of c-quark in the $|uudc\bar{c}\rangle$ state: in the intrinsic charm model [1] satisfied (5) (solid line); in the alternative approach based on [4] model satisfied (9) (dashed line)

As we can see in Fig. 1 the probability distributions (5) and (9) are almost the same, under paremeters we chose. So we can suppose that the presented in ref. [1] probability distribution refers to the c-quark distribution in the pentaquark $|uudc\bar{c}\rangle$ and was nothing to do with sea c-quarks.

We have one more argument in favor of the hypothesis. It is well known that the quark distribution functions in K-mesons and π-mesons are different. The diffrence can be explained by the model (7) (see [7]) on the one hand and by (1) model on the other hand. It convinces us that the probability distributions (5) and (9) refer to the valence c-quark momentum distribution in the pentaquark $|uudc\bar{c}\rangle$.

The next problem appeared is the proton-pentaquark mixing which determs absolute normalization. Let matrix element for $p \to |uudc\bar{c}\rangle$ be equal to V, then admixture of the pentaquark in a proton is given by factor $V/(E_1 - E_2)$, where E_1, E_2 are energies of the pentaquark and the nucleon respectively. The matrix element can be evaluated as Λ_{QCD} and the energy diffrence can be estimated as the diffrence of masses. In that way the posibility to find the pentaquark in a proton is less than:

$$W(p \to uudc\bar{c}) \leq \left(\frac{300MeV}{3GeV}\right)^2 \approx 1\% \qquad (10)$$

In fact we should expect the mixing to be much smaller. Most likely it is due to diffrent color state of quarks in the pentequark and in a proton.

1. In the original work [1], based on assumption that most of the charm cross-section comes from diffraction in pp-interaction, the mixing was evaluated by 1%.

2. In the later work [8] the restriction 0.59% was given, based on data of European Muon Collaboration (EMC) in hadronic scattering. The best fit to data gave 0.31%.

3. Differential spectrum of D-mesons in photoproduction processes [2] contradicts the intrinsic charm hypothesis and set the contribution restriction in the range of $\sim 0.1 - 0.2\%$.

4. The most fruitful results concerning a test of the mixing hysothesis can be obtained by hidden charm particles production. Such analysis was performed in ref. [3]. The limitation for the fluctuation $p \to |uudc\bar{c}\rangle$ probability is $\sim 10^{-7}$, according to [3].

The discussed model, based on the assumption about the pentaquark-proton mixing, gives a consistent interpretation of the intrinsic charm phenomenon, which is used so far for the hadron physics application [9]. It is worth emphesizing that the observation of the intrinsic charm phenomenon directly results in the existance of the stable pentaquark $|uudc\bar{c}\rangle$. A discovery of pentaquark $|uudc\bar{c}\rangle$ or getting more stringent restriction is a task of future experiments.

Author would like to thank Prof. A.K.Likhoded for fruitful discussions and a set of constructive remarks.

References

[1] *S.J. Brodsky, P.Hoyer, C.Peterson, N.Sakai*, Phys. Lett., **93B** (1980), 451.

[2] *S.S. Gershtein, A.K. Likhoded, S.R. Slabospitsky*, Nucl. Phys.., **34** (1981), 227.

[3] *A.K. Likhoded, V.A. Litvin*, Nucl. Phys.., **62** (1999), 728-741.

[4] *J. Kuti, V.F. Veisskopf*, Phys. Rev. **D4** (1971) 3418.

[5] *V.G. Kartvelishvili, A.K. Likhoded, V.A. Petrov*, Phys. Lett., **78B** (1978), 615.

[6] *S.Drell, T.M.Yan* Phys. Rev. **24** (1970), 181.
G.B.West Phys. Rev. **24** (1970), 1206.

[7] *P.V. Chliapnikov, V.G. Kartvelishvili, V.V.Kniasev, A.K. Likhoded*, Nucl. Phys.., **B148** (1979), 400-412.

[8] *E. Hoffmanm, R. Moore*, Z. Phys. **C20** (1983), 71

[9] *G.I.Lykasov, V.A.Bednyakov, A.F.Pikelner, N.I.Zimin*, arXiv:**1205.1131v2** (2012).